

AD-753 442

MILOCMED 1968: A DROGUE EXPERIMENT IN
THE IONIAN SEA. PART I. THE TEMPERATURE
TIME SERIES

Adolf Dahme, et al

SACLANT ASW Research Centre
La Spezia, Italy

15 October 1972

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE
5285 Port Royal Road, Springfield Va. 22151

AD 753442




SACLANT ASW
RESEARCH CENTRE

SACLANTCEN
Technical Memorandum No. 181

MILOMED 1968: A DROGUE EXPERIMENT IN THE IONIAN SEA
PART I: THE TEMPERATURE TIME SERIES

by

ADOLF DAHME and ANDRE DE HAEN



15 OCTOBER 1972

This document has been approved
for publication and sale, its
distribution is unlimited.

NORTH
ATLANTIC
TREATY
ORGANIZATION

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
U.S. Department of Commerce
Washington, D.C. 20540

VIALE SAN BARTOLOMEO 400
I-19026 - LA SPEZIA, ITALY

This document is unclassified. However, the information it contains is published subject to the conditions of the legend printed on the inside cover. Short quotations from it may be made in other scientific publications if credit is given to the author(s) and to SACLANTCEN, requests for other reproduction, except in official NATO publications, should be addressed to the Director, SACLANTCEN.

1. The recipient NATO Government agrees to use its best endeavours to ensure that the information herein disclosed, whether or not it bears a security classification, is not dealt with in any manner (a) contrary to the intent of the provisions of the Charter of the Centre, or (b) prejudicial to the rights of the owner thereof to obtain patent, copyright, or other like statutory protection therefor.

ADDRESS ONLY

NAME

DATE

TIME

BY

RECEIVED

1966

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

SACLANTCEN
TECHNICAL MEMORANDUM NO. 181

NORTH ATLANTIC TREATY ORGANIZATION
SACLANT ASW RESEARCH CENTRE
Viale San Bartolomeo 400
I 19026 - La Spezia, Italy


MILOCMED 1968: A DROGUE EXPERIMENT IN THE IONIAN SEA
PART I: THE TEMPERATURE TIME SERIES

by
Adolf Dahme and André de Haen

Published

15 October 1972

APPROVED FOR DISTRIBUTION
(June 1972)



Ir M.W. van Batenburg
Director

Manuscript Completed:

15 March 1971

I 2

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	1
INTRODUCTION	2
1. DATA COLLECTION	3
2. DATA PROCESSING	8
3. RESULTS	10
3.1 Results Derived from Daily Means	10
3.2 Results Derived from Hourly Means	13
CONCLUSIONS	31
REFERENCES	33

List of Figures

1. Type of drogue used in MILOCMED 68	6
2. Track chart of drogue alpha and drogue foxtrot	7
3a. Daily mean temperatures at selected depths: drogue alpha	20
3b. Daily mean temperatures in the upper 15 m: drogue alpha	21
3c. Standard deviations of temperatures in the upper 10 m (1-day periods): drogue alpha	21
4a. Daily mean temperatures at selected depths: drogue foxtrot	22
4b. Daily mean temperatures in the upper 15 m: drogue foxtrot	22
4c. Standard deviations of temperatures in the upper 10 m: (1-day periods): drogue foxtrot	22
5. Temperature profiles 25-28 May 1968, based on daily means recorded by drogue alpha	23
6a. Temperature profiles of drogue alpha and drogue foxtrot, 17 May 1968	24
6b. Temperature profiles of drogue alpha and drogue foxtrot, 18 May 1968	24
6c. Temperature profiles of drogue alpha and drogue foxtrot, 19 May 1968	24
6d. Temperature profiles for drogue alpha on 18 May and drogue foxtrot on 17 May 1968	25
6e. Temperature profiles for drogue alpha on 19 May and drogue foxtrot on 18 May 1968	25
6f. Temperature profiles of drogue alpha for 18 and 19 May 1968	26
6g. Temperature profiles of drogue foxtrot for 18 and 19 May 1968	26

TABLE OF CONTENTS (Cont'd)

List of Figures (Cont'd)

	<u>Page</u>
7. Differences in temperature between drogue alpha and drogue foxtrot as a function of their distances apart	27
8. Hourly mean temperatures and standard deviations computed from the records of drogue alpha	28
9. Hourly mean temperatures and standard deviations computed from the records of drogue foxtrot	29
10. Heat content (in excess of 17°C) in calories per unit cross section of the total water column above indicated depths; drogue alpha	30

MILOCMED 1968: A DROGUE EXPERIMENT IN THE IONIAN SEA

PART I: THE TEMPERATURE TIME SERIES

by

Adolf Dahme and André de Haen

ABSTRACT

During an oceanographic survey (MILOCMED 68) made in the Ionian Sea between 15 May and 5 June 1968, temperature profiles were obtained every three minutes by means of free-drifting, self-recording drogues equipped with 60 m long thermistor chains. The gathering and processing of two of these time series, one covering a period of 17 days and one a period of 4 days, are described. The results reported include an indication of an oceanographic front in the area that causes sharp horizontal changes in the temperature structure both above and below the thermocline. They also include information about the daily heating cycle of the near-surface water and about the effects of internal waves on the temperature structure within the thermocline.

INTRODUCTION

An oceanographic survey of an area approximately 100 miles east of Malta was made between 15 May and 5 June 1968 (MILOCMED 68) with the assistance of several nations. Part of the programme was a study of the diffusion of the surface water, made by tracking the drift of six drogues, which is described in a companion report (Ref. 1).

The present paper describes the temperature variations observed by arrays of 21 thermistors suspended beneath two of the drogues (ALPHA, FOXTROT) at fixed depths between 0.5 m and 60 m. (Unfortunately, the thermistor arrays attached to two other drogues malfunctioned and the data could not be meaningfully processed.) The period covered by drogue ALPHA extends from 1615Z 16 May to 1400Z 2 June and by drogue FOXTROT from 1448Z 16 May to 1515Z 20 May.

This report describes in some detail the techniques employed for the collection and processing of the temperature profile data; the main part of the report describes several remarkable temperature variations recorded as the drogues drifted through different water masses.

1. DATA COLLECTION

Figure 1 gives a simplified sketch of the drogues used to support the thermistor arrays. Each drogue had four 5 m² canvas-covered vanes (mounted axially at right angles to each other) to increase drag. A mast about 3 m high carried a radar reflector and a flashing light to facilitate tracking.

All the thermistors were electrically connected to outlets moulded at specified positions [see Table 1] on neoprene-insulated, concentric twin cable of 9 mm diameter. For the thermistors suspended below the drogue the cable was taped to a 3.2 mm diameter wire rope carrying a 60 kg weight. The cable for the shallower thermistors was taped to a nylon rope stretched along the side of the drogue. The error in geometrical distance of the thermistors from the design datum level (the sea surface) is believed to be less than ± 5 cm and, as the drogues had enough buoyancy to ride any waves more than a few metres long, this distance can be considered as the true depth of the thermistors beneath the free sea surface.

Each thermistor was housed in a small plastic case to increase its time constant to about two minutes, that is, $1/3$ of the Nyquist period based on the sampling period of three minutes. This increased time constant also has the advantage of reducing the influence of wave motion, as it is much larger than the wave periods. Thus the variations in the temperature structure are integrated by the fact that the array closely follows the wave motion.

The scanning and recording procedures performed by equipment housed in the cylindrical body of the drogues were essentially those described in Ref. 2 and successfully used during previous surveys in the Strait of Gibraltar. All thermistors were scanned every three minutes, one whole scan being completed in 10 seconds.

The signals were recorded on tape for later processing by the computer ashore. An additional feature not available during previous cruises was that a small transmitter housed in the upper part of the drogue could transmit the signals present on the tape over distances of a few miles, thereby indicating whether the whole temperature recording system was working properly. The absolute accuracy of the complete system, including calibration, recording, computer transfer and processing, is estimated to be 0.1°C , but the relative accuracy between adjacent scans is probably better than 0.02°C .

The tracks of the two drogues while useful data were being recorded are given in Fig. 2. Drogue ALPHA was launched at 1554Z on 16 May and by 1615Z its array started to make undisturbed temperature recordings. The drogue was partly lifted between 0900Z and 0954Z on 20 May for inspection, without changing its position or interrupting the electronic circuitry. At 0430Z on 23 May it was recovered for repairs to the canvas of the vanes. When it was relaunched on the same day at 0742Z it was placed approximately 10 n.mi west of its previous position so as to make it easier to track the two drogues, which had by then separated considerably. There were no further interruptions until it was recovered at 0800Z on 5 June. However, inspection of the data revealed that the battery power had dropped during the last days, causing a drift of the temperature data. Hence, only the data recorded until 1400Z on 2 June were considered good.

Drogue FOXTROT was launched approximately 5 n.mi south of Drogue ALPHA at 1348Z on 16 May and undisturbed recording started at 1448Z. The drogue was recovered for repair at 1518Z on 20 May. After relaunching at 1827Z on 20 May the drogue ceased to give useful data.

TABLE 1

DEPTHS OF THERMISTORS

DROGUE ALPHA

0.50 m
1.25 m
2.25 m
3.25 m
4.00 m
5.00 m
6.00 m
7.00 m
8.00 m
9.00 m
10.00 m
12.00 m
14.00 m
16.00 m
18.00 m
22.00 m
28.00 m
34.00 m
40.00 m
50.00 m
60.00 m

DROGUE FOXTROT

0.50 m
1.25 m
2.00 m
2.75 m
3.25 m
4.00 m
5.00 m
6.00 m
7.00 m
8.00 m
9.00 m
10.00 m
12.00 m
14.00 m
18.00 m
22.00 m
28.00 m
34.00 m
40.00 m
50.00 m
60.00 m

DROGUE MILOC 1968




FIG. 1 TYPE OF DROGUE USED IN MILOCMED 68




FIG. 2 TRACK CHART OF DROGUE ALPHA AND DROGUE FOXTROT
(The track of Drogue ALPHA during 26-28 May is repeated in bottom left corner)

2. DATA PROCESSING

Before the recordings were converted from analogue to digital form and transferred to the computer, their quality was checked by displaying the data on an electrosensitive strip chart in the same form as it had been recorded on magnetic tape. This also permitted a check of the start and end of the undisturbed recording, the noise level, the regularity of the synchronization pulses, and the time marks.

The data were then processed by a general computer system for oceanographic data developed at SACLANTCEN, out of which the following sets of programs were used:

One program checks the homogeneity of the calibration points of the thermistor probes and calculates their deviation from the theoretical experimental curves; after any necessary adjustments another program calculates the coefficients of the functions representing the individual calibration curves.

The asynchronously-stored time intervals resulting from the transfer from the tapes are separated into individual scans. The successive time intervals are then referred to a common origin and assigned to their corresponding sensors. One auxiliary program checks this synchronization and another counts the cycles found between two fixed time marks in order to make sure that no scan has been lost, which would result in a shifted time scale for the rest of the analysis. The response times of the sensors are then introduced into their respective calibration equations and the corresponding temperatures are calculated.

The original recordings contain reference information that is transferred to successive computer tapes and used to detect incomplete, noisy or unsynchronized scans, which would result in errors in the calculated temperatures. If such a fault is present,

part of the scan is usually incorrect. This part is then rejected by the program and replaced by a linear interpolation between the preceding and subsequent accepted values. These eliminated scans represented less than 1% of the total data reported here. The original record has a time mark instead of a temperature profile at every 80th scan, i.e. every 4 hours of real time; this missing scan is also replaced by an interpolation. To avoid an interruption of the time series the two large gaps (one of 1 hour and one of 3 hours) in the data from drogue ALPHA were filled by a series of temperature profiles linearly interpolated between the preceding and subsequent good scans.

The refined data were finally treated by specific programs that compute, list and plot the hourly and daily means and their corresponding standard deviations, as well as computing the total heat content of selected water columns and plotting the results against time.

3. RESULTS

3.1 Results derived from Daily Means

Tables 2 and 3 list the means and standard deviations of the 480 measurements recorded each day by drogues ALPHA and FOXTROT respectively. Because of the behaviour of the standard deviation — borne out by the hourly means — for the 3.25 m level of Drogue ALPHA, the values for this level from 17 to 26 May (given in parenthesis in Table 2) should be ignored.

The changes in the temperature structure during the survey can be seen more clearly from Figs. 3a and 4a, which display the daily mean temperatures at depths from 10 m to 60 m by 10 m increments (the levels at 20 m and 30 m are linearly interpolated from the adjacent level). The connecting lines serve solely to assist the eye to identify values belonging to the same level.

3.1.1 Drogue ALPHA

The most striking feature in the records of the mean temperature structure obtained from this drogue is the drastic changes between 26 and 28 May. In this period the temperature of the water at and below 20 m increased by about 2°C in the upper part of this stratum and by about 1°C in its lower part. This is demonstrated in a more familiar way in Fig. 5, which shows the temperature profiles in the form of bathythermograms. At the same time the temperature at the 10 m level decreased by nearly 1°C, partly through wind mixing. Unfortunately, the absence of salinity records for this period prevents a study of the change in the density structure and of the possible change in stability. The drogue's track on 26 and 27 May, which is repeated in the bottom left-hand corner of Fig. 2 for increased clarity, does not show any peculiarity that might help to explain this drastic change of the temperature structure, nor do the meteorological observations made aboard MARIA PAOLINA and

PLANET give a hint. At the end of the survey period the temperature tends to return to values obtained during the first part of the survey.

An idea of the mean temperature gradient in the surface layer above the thermocline is given by Fig. 3b, which shows the daily mean temperatures at 0.5 m, 5 m, 10 m and 15 m (the last being linearly interpolated from the 14 m and 16 m levels). During the first nine days there was a general heating at these levels, associated with an increasing temperature gradient. An exception was observed at the 16 m level on 20 May when it was apparently affected by an exceptionally large internal wave. The drastic changes at the lower levels on 27 and 28 May are reflected at the upper levels, but in the opposite sense: the temperature at 0.5 m depth decreased during these days by approximately 2°C. On these and the following two days the water at the four levels shown in Fig. 3b appears well mixed, which is confirmed by the standard deviations of the temperatures at the 0.5 m, 5 m and 10 m levels recorded on Fig. 3c. A considerable temperature gradient again developed on 1 June.

In Table 2b, the maximum and minimum values of the standard deviations of the temperatures are emphasized by circles. The depth of the maximum value, approximately indicating the depth where internal waves are most active (in as much as they produce the largest temperature variations) is constantly at 18 m until 26 May, and descends to 40 m during 27 May, concurrently with the change observed in Fig. 3a. It then ascends to 22 m during the last three days. The depth of the minimum standard deviation is more variable, the biggest change again occurring between 26 and 27 May. As previously indicated, the standard deviation is also a useful tool for investigating how homogeneous the different levels were or how well mixed the layer above the thermocline was; on 28 May, for example, the layer was particularly well mixed.

3.1.2 Drogue FOXTROT

Unfortunately the records of Drogue FOXTROT are short. They will therefore be used mainly for a study of the differences in the temperature structure at the different positions of the two drogues during the period of 17-19 May.

3.1.3 Differences of the Temperature Structure at the Positions of Drogue ALPHA and Drogue FOXTROT

Figures 6a, b and c compare the temperature structures (based on daily means) at the positions of the two drogues on 17, 18 and 19 May respectively and Table 4 lists the actual differences in temperature at each depth.

It is seen that, whereas there was little difference between the temperature profiles on 17 May [Fig. 6a] — in fact they were remarkably similar at depths below 25 m — there were differences on 18 and 19 May [Figs. 6b and 6c]. One is tempted to assign these differences to the increasing distance between the two drogues, as recorded in Table 5, which lists the distances every two hours (taken from the track chart of Fig. 2) together with the calculated daily average. At levels within the thermocline the temperature differences between the two drogues cannot properly be compared, because here the temperature gradients are stronger and therefore the influence of internal waves appears greater. In the survey area this might be assumed to occur at the four depths of 14 m, 18 m, 22 m and 28 m marked on Table 4. Above the thermocline, the picture is not uniform, as is demonstrated by Figs. 7a and 7b, based on Tables 4 and 5. For depths between the surface and 7 m, Fig. 7a suggests that on 17 and 18 May the two drogues stayed in more or less the same water mass but that on 19 May one drogue passed a boundary or a front. At greater depths, but still above the thermocline [Fig. 7b], no certain conclusion can be drawn. Below the thermocline [Fig. 7c] it appears that a front may have appeared between the two drogues on 18 May and remained there on 19 May.

Taking the geographical positions, and not the distances, as the determining factors, one observes from the track chart [Fig. 2] that nearly the same area was crossed by Drogue FOXTROT on 17 May as by Drogue ALPHA on 18 May. The same happened on the following days: an area crossed by Drogue FOXTROT on 18 May was crossed by Drogue ALPHA on 19 May. The temperature profiles for these two areas are plotted in Figs. 6d and 6e and it is seen that in both areas the temperature structure changed from one day to the other. However, as revealed by Figs. 6f and 6g, neither drogue recorded a change in the temperature structure beneath the thermocline on the two consecutive days. (Because the temperature structures above the thermocline are also influenced by the changing meteorological conditions they cannot be directly compared.) It can therefore be concluded that the boundary between the two water masses located beneath the thermocline and between the two drogues moved eastwards during 18 and 19 May.

It is noted that when Drogue ALPHA passed between 0 and 4 n.mi south of the same area on 27 and 28 May the water beneath the thermocline was considerably warmer [Fig. 3a].

3.2 Results Derived from Hourly Means

Figures 8 and 9 present graphically, as a function of time, the means and standard deviations of the 20 measurements recorded each hour by Drogues ALPHA and FOXTROT respectively.

The daily heating cycle within the surface layer is clearly demonstrated, its effect being normally confined to about the upper 8 m. Only when strong winds prevailed throughout the day, as on 20, 21 and 20 May, was the penetration deeper, whereas on calm days the penetration was less. The sharp peaks in the very near-surface layer on 25 May and 1 June were associated with calm weather at midday. The drastic changes on 26 and 27 May are easily recognizable: a decrease of the temperature in the surface layer is accompanied by an increase of the temperature below the

thermocline. At the same time the "layer depth", i.e. the top of the thermocline, increases from about 14 m or 15 m to more than 18 m.

The strong oscillations of the temperature around depths of 22 m are attributed to internal waves. The thermal unrest extends, but to a lesser degree, to the region below the thermocline as a consequence of the temperature gradients still present in this stratum. Note that no changes in the temperature structure occurred when Drogue ALPHA was relaunched on the morning of 23 May at a position 10 n.mi west of its previous position.

Figure 10 shows the heat content, H , in excess of 17°C of a water column of 1 cm^2 cross-section between the surface and the indicated depths Z_c , computed as

$$H = \rho_0 c_p \int_0^{Z_c} (T - 17) dz,$$

where

ρ_0 = specific gravity of the water.

c_p = specific heat of the water under constant pressure.

T = temperature at depth z .

The general heating occurring during the periods 17 to 25 May (the first part of the 0-14 m curve indicates a daily average net heat gain of 200 cal/cm^2 for the 14 m deep water column during the first ten days of the survey) and 28 May to 1 June is clearly seen. On 26 and 27 May the drogue entered cooler surface water and there was a great apparent loss of heat. Another loss of heat started on 2 June.

TABLE 2a

Daily Mean Temperature (°C) Computed from the 7 records of Bregu ALPHA

	May 1965																June
Depth m	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	
0.50	19.997	20.117	20.118	20.156	20.173	20.180	20.936	21.980	21.973	21.687	20.878	20.322	20.197	20.100	20.090	21.256	
1.25	19.975	20.119	20.080	20.101	20.174	20.737	20.829	21.116	21.736	21.831	20.681	20.317	20.190	20.100	20.079	21.107	
2.25	19.993	20.119	20.081	20.106	20.177	20.739	20.809	21.114	21.689	21.687	20.677	20.305	20.190	20.700	20.070	21.034	
3.25	(19.985)	(20.123)	(20.107)	(20.120)	(20.174)	(20.660)	(20.611)	(20.933)	(21.252)	(21.333)	20.606	20.100	20.012	20.119	20.098	21.063	
4.00	19.952	20.016	20.130	20.133	20.160	20.700	20.832	20.990	21.435	21.685	20.707	20.321	20.623	20.124	21.003	21.054	
5.00	19.955	20.110	20.120	20.122	20.160	20.115	20.793	20.911	21.237	21.650	20.730	20.327	20.623	20.111	20.697	20.080	
6.00	19.960	20.095	20.100	20.095	20.130	20.800	20.731	20.979	21.024	21.704	20.606	20.300	20.600	20.770	21.054	20.012	
7.00	19.957	20.093	20.101	20.070	20.110	20.884	20.632	20.963	20.934	21.323	20.605	20.270	20.570	20.740	20.912	20.450	
8.00	19.905	20.071	20.110	20.101	20.097	20.863	20.660	20.967	20.910	21.253	20.601	20.303	20.600	20.791	20.923	20.970	
9.00	19.957	20.060	20.112	20.100	20.097	20.860	20.674	20.963	20.951	21.220	20.600	20.304	20.620	20.700	20.610	20.865	
10.00	19.950	20.055	20.107	20.100	20.091	20.855	20.650	20.950	20.944	21.170	20.607	20.301	20.613	20.795	20.904	20.966	
12.00	19.964	20.060	20.073	19.995	20.173	20.714	20.104	20.755	20.722	20.955	20.612	20.223	20.535	20.725	20.705	20.755	
14.00	19.975	20.015	20.000	19.960	20.201	20.770	20.315	20.832	20.693	20.912	20.616	20.260	20.570	20.773	20.922	20.750	
16.00	19.918	20.063	19.925	19.937	19.912	19.910	19.970	20.110	20.773	20.653	20.648	20.174	20.606	20.660	20.660	20.582	
18.00	19.918	19.996	19.991	19.997	19.921	19.791	19.910	19.970	19.935	19.604	20.378	20.220	20.523	20.605	20.605	20.604	
22.00	17.210	17.150	17.755	17.620	17.321	17.618	17.650	17.690	17.602	19.223	19.510	20.100	20.624	20.030	20.144	19.212	
24.00	16.910	17.033	17.000	16.906	16.710	16.937	16.932	16.961	16.964	17.137	19.105	19.635	19.607	19.605	19.605	17.574	
34.00	16.510	16.731	16.715	16.694	16.335	16.606	16.660	16.701	16.655	16.605	17.106	16.100	17.610	17.567	17.511	16.916	
40.00	16.100	16.305	16.165	16.090	15.988	15.975	16.042	15.954	15.973	16.007	16.507	17.644	17.620	17.654	17.234	16.800	
50.00	15.627	15.905	15.900	15.547	15.458	15.404	15.547	15.671	15.673	16.157	16.157	16.400	16.505	16.484	16.051	15.953	
60.00	15.301	15.517	15.513	15.205	15.101	15.172	15.225	15.164	15.194	15.277	15.967	16.175	15.911	15.951	16.053	15.521	

TABLE 2b

Standard Deviations (°C) of the Temperatures Recorded by Bregu ALPHA for Periods of One Day

	May 1965																June
Depth m	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	1	
0.50	0.067	0.124	0.234	0.174	0.131	0.268	0.254	0.500	0.350	0.113	0.200	0.078	0.191	0.135	0.142	0.450	
1.25	0.067	0.125	0.224	0.173	0.120	0.255	0.251	0.550	0.340	0.113	0.200	0.078	0.191	0.133	0.130	0.217	
2.25	0.060	0.124	0.221	0.172	0.125	0.255	0.245	0.354	0.340	0.116	0.200	0.074	0.191	0.132	0.135	0.166	
3.25	(0.166)	(0.132)	(0.223)	(0.115)	(0.137)	(0.375)	(0.445)	(0.523)	(0.603)	(0.406)	0.290	0.077	0.191	0.132	0.135	0.163	
4.00	0.060	0.124	0.207	(0.174)	0.126	0.250	0.137	0.224	0.414	0.170	0.300	0.070	0.191	0.130	0.120	0.136	
5.00	0.060	0.115	0.172	0.174	0.111	0.190	0.089	0.112	0.343	(0.118)	0.207	0.070	0.191	0.121	0.111	0.114	
6.00	0.050	0.116	0.137	0.152	0.067	0.107	0.057	0.070	0.204	0.135	0.207	0.074	0.190	0.104	0.105	0.070	
7.00	0.055	0.102	0.102	0.201	0.052	0.040	0.002	0.066	0.110	0.142	0.206	0.070	0.190	0.004	0.050	0.047	
8.00	0.055	0.097	0.060	0.214	0.071	0.026	0.001	0.055	0.057	0.201	0.205	0.066	0.190	0.000	0.001	0.035	
9.00	(0.154)	0.090	0.039	0.204	(0.061)	0.023	0.055	0.056	(0.055)	0.220	0.202	0.065	0.190	0.050	0.044	(0.033)	
10.00	0.054	0.057	(0.028)	0.208	0.003	0.038	(0.057)	(0.056)	0.001	0.244	0.207	0.060	0.190	0.045	0.035	0.030	
12.00	(0.051)	(0.070)	0.055	0.266	0.054	(0.010)	0.213	0.058	0.138	0.242	0.201	0.056	0.192	0.042	(0.037)	0.050	
14.00	0.166	0.050	0.145	0.303	0.126	0.071	0.454	0.517	0.406	0.275	0.261	0.050	0.177	(0.051)	0.064	0.140	
16.00	0.173	0.314	0.313	0.310	0.333	0.506	0.441	0.567	0.308	0.372	(0.280)	(0.055)	0.144	0.000	0.006	0.127	
18.00	(0.457)	(0.782)	(0.100)	(0.500)	(0.716)	(0.912)	(0.655)	(0.937)	(0.558)	(1.004)	0.280	0.055	(0.110)	0.210	0.102	0.325	
22.00	0.170	0.473	0.145	0.315	0.236	0.225	0.268	0.244	0.519	0.885	0.402	0.160	0.444	(0.470)	(0.461)	(0.703)	
24.00	0.115	0.171	0.151	0.115	0.160	0.156	0.115	0.197	0.207	0.270	0.586	(0.338)	(0.665)	0.436	0.426	0.343	
34.00	0.114	0.110	0.142	0.151	0.174	0.140	0.110	0.170	0.167	0.241	0.717	0.280	0.586	0.260	0.251	0.302	
40.00	0.121	0.161	0.153	0.106	0.164	0.144	0.120	0.140	0.153	0.274	(0.575)	0.157	0.405	0.222	0.240	0.104	
50.00	0.101	0.114	0.160	0.144	0.066	0.063	0.111	0.083	0.071	0.166	0.567	0.125	0.256	0.206	0.171	0.160	
60.00	0.115	0.105	0.115	0.132	0.054	0.057	0.060	0.060	0.054	0.153	0.242	0.105	0.144	0.147	0.205	0.104	

TABLE 3a

Daily Mean Temperatures (°C)
as Computed from the records of Drogue FOXTROT

Depth m	17	18	19
0.50	20.018	20.124	20.308
1.25	19.960	20.042	20.231
2.00	19.965	20.069	20.228
3.25	19.957	20.056	20.211
4.00	19.982	20.082	20.226
5.00	19.930	20.014	20.105
6.00	19.947	20.030	20.051
7.00	19.887	19.982	19.942
8.00	19.954	20.171	20.178
9.00	19.788	19.934	19.881
10.00	19.730	19.894	19.868
12.00	19.638	19.800	19.868
14.00	19.571	19.706	19.830
18.00	18.809	18.878	19.318
22.00	17.550	17.527	18.380
28.00	16.833	16.875	16.986
34.00	16.513	16.501	16.537
40.00	16.114	16.013	16.037
50.00	15.626	15.486	15.498
60.00	15.365	15.210	15.206

TABLE 3b

Standard Deviations (°C) of the Temperatures
Recorded by Drogue FOXTROT for Periods of One Day

Depth m	17	18	19
0.50	0.118	0.149	0.264
1.25	0.115	0.414	0.259
2.00	0.127	0.172	0.264
3.25	0.126	0.143	0.249
4.00	0.116	0.143	0.246
5.00	0.112	0.125	0.227
6.00	0.114	0.115	0.189
7.00	0.129	0.104	0.121
8.00	0.191	0.105	0.053
9.00	0.181	0.073	0.023
10.00	0.178	0.072	0.019
12.00	0.136	0.119	0.048
14.00	0.192	0.184	0.151
18.00	0.564	0.618	0.331
22.00	0.420	0.412	0.702
28.00	0.201	0.170	0.397
34.00	0.143	0.142	0.238
40.00	0.137	0.130	0.189
50.00	0.155	0.156	0.123
60.00	0.126	0.138	0.108

TABLE 4

Differences ($^{\circ}\text{C}$) Between the Temperature Recorded at Drogue ALPHA
and the Temperature at Drogue FOXTROT for the Same Depths

Depth m	17	18	19
0.50	- 0.051	- 0.011	+ 0.106
1.25	+ 0.018	+ 0.072	+ 0.178
4.00	- 0.030	+ 0.016	+ 0.110
5.00	+ 0.055	+ 0.102	+ 0.175
6.00	+ 0.013	+ 0.055	+ 0.145
7.00	+ 0.100	+ 0.111	+ 0.219
8.00	+ 0.011	- 0.100	- 0.048
9.00	+ 0.169	+ 0.126	+ 0.231
10.00	+ 0.229	+ 0.164	+ 0.239
12.00	+ 0.306	+ 0.240	+ 0.205
14.00	+ 0.304	+ 0.312	+ 0.178
18.00	- 0.495	+ 0.526	+ 0.073
22.00	- 0.334	+ 0.129	- 0.625
28.00	- 0.023	+ 0.158	+ 0.054
34.00	+ 0.003	+ 0.230	+ 0.178
40.00	+ 0.032	+ 0.352	+ 0.311
50.00	+ 0.001	+ 0.412	+ 0.362
60.00	+ 0.026	+ 0.307	+ 0.307

} thermocline

TABLE 5

Distances apart (n.mi) of Droque ALPHA and Droque FOXTROT
at different times

Hour GMT	16	17	18	19	20
0	-	6.3	8.3	11.1	10.9
2	-	5.6	9.1	11.1	10.7
4	-	5.6	9.1	10.7	10.7
6	-	5.5	9.7	10.6	10.0
8	-	5.2	9.3	10.7	7.9
10	-	6.7	9.8	11.2	10.7
12	-	6.7	10.1	10.7	11.0
14	-	6.6	9.3	11.4	11.0
16	6.5	7.0	8.7	11.4	-
18	4.6	7.1	10.4	10.6	-
20	5.5	7.3	9.6	10.6	-
22	6.3	7.5	10.3	10.6	-
mean	-	6.3	9.5	10.9	-




FIG. 3a DAILY MEAN TEMPERATURES AT SELECTED DEPTHS: DROGUE ALPHA




FIG. 3b DAILY MEAN TEMPERATURES IN THE UPPER 15 m: DROGUE ALPHA




FIG. 3c STANDARD DEVIATIONS OF TEMPERATURES IN THE UPPER 10 m (1-day periods): DROGUE ALPHA




FIG. 4a DAILY MEAN TEMPERATURES AT SELECTED DEPTHS: DROGUE FOXTROT




FIG. 4b DAILY MEAN TEMPERATURES IN THE UPPER 15 m: DROGUE FOXTROT




FIG. 4c STANDARD DEVIATIONS OF TEMPERATURES IN THE UPPER 10 m (1-day periods): DROGUE FOXTROT




FIG. 5 TEMPERATURE PROFILES 25-28 MAY 1968, BASED ON DAILY MEANS RECORDED BY DROGUE ALPHA




FIG. 6a. TEMPERATURE PROFILES OF DROGUE ALPHA AND DROGUE FOXTROT, 17 MAY 1968

24.1




FIG. 6b. TEMPERATURE PROFILES OF DROGUE ALPHA AND DROGUE FOXTROT, 18 MAY 1968

24.2




FIG. 6c. TEMPERATURE PROFILES OF DROGUE ALPHA AND DROGUE FOXTROT, 19 MAY 1968

24.3




FIG. 6 TEMPERATURE PROFILES FOR DROGUE ALPHA ON 18 MAY AND DROGUE FOXTROT ON 17 MAY 1968 (approximately the same track)




FIG. 6e TEMPERATURE PROFILES FOR DROGUE ALPHA ON 19 MAY AND DROGUE FOXTROT ON 18 MAY 1968 (approximately the same track)




FIG. 6f TEMPERATURE PROFILES OF DROGUE ALPHA FOR 18 AND 19 MAY 1968




FIG. 6g TEMPERATURE PROFILES OF DROGUE FOXTROT FOR 18 AND 19 MAY 1968





FIG. 7 DIFFERENCES IN TEMPERATURE BETWEEN DROGUE ALPHA AND DROGUE FOXTROT AS A FUNCTION OF THEIR DISTANCES APART
a. 0.5 m to 7 m
b. 8 m to 12 m
c. 34 m to 60 m

27.2

27.3

MILOC MED 68

DROGUE ALPHA



Reproduced from
best available copy.

FIG. 8 HOURLY MEAN TEMPERATURES AND STANDARD DEVIATIONS COMPUTED FROM THE RECORDS OF DROGUE ALPHA (thick lines are hourly means, fine lines are standard deviations)




FIG. 9 HOURLY MEAN TEMPERATURES AND STANDARD DEVIATIONS COMPUTED FROM THE RECORDS OF DROGUE FOXTROT (thick lines are hourly means, fine lines are standard deviations)

MILOC MED 68 DROGUE ALPHA




FIG. 10 HEAT CONTENT (IN EXCESS OF 17°C) IN CALORIES PER UNIT CROSS SECTION OF THE TOTAL WATER COLUMN ABOVE INDICATED DEPTHS; DROGUE ALPHA

CONCLUSIONS

So far only a very limited analysis of the data has been performed; its results may be summarized as follows:

(1) During the heating season a very pronounced daily heating cycle is observed in the upper waters of the Ionian Sea, but this variation is limited in depth to eight metres except on days with prevailing strong winds. On the other hand, calm days cause a sharp increase of the temperature very close to the surface, but there is no longer a daily heating cycle throughout the upper eight metres. There is no reason why this result should not be extended to the whole Mediterranean.

(2) Internal waves cause the temperature structure within the thermocline to change very rapidly. Although no spectral analysis has been done, the high standard deviations found for 1-hour periods suggest that the periods of these temperature variations are less than one hour.

(3) The data support the concept that horizontal temperature changes — above and below the thermocline — are not steady but occur at fairly sharp boundaries. In other words, water bodies of unknown shape and size are moving through the ocean and taking a very long time to lose their identity by mixing in the horizontal direction. It is recommended that the size of these bodies should be investigated in more detail.

Since the manuscript of this report was written there has been a further analysis of MILOC 68 data by P. Saunders of Woods Hole and F. Edwards (Royal Navy attached to SACLANTCEN). Further cruises have been made and the analysis of the results [some published (Refs. 3, 4) and some unpublished] by Woods of the UK Meteorological Office and by Briscoe and Johannessen of SACLANTCEN confirm the existence of an oceanic front or system of fronts in the area.

In particular, this frontal system appears to extend into comparatively shallow water east of Malta in summer and late spring and has been detected in deep water approximately 100 n.mi east of Malta in December. Unfortunately the drogue data reported here lack sufficient information on the lateral extent of the large variations recorded on 26-28 May to "prove" that a front was also present in the deep water in late spring — but it is an intriguing thought.

REFERENCES

1. Molcard, R. 1972. "MEDMILOC 68/Drogue Experiment Part II: A Study of a Large-Scale Diffusion", SACLANTCEN Technical Memorandum No. 182.
2. Ziegenbein, J. 1969. "The Gibraltar Buoy System", SACLANTCEN Technical Memorandum No. 144.

[AD 856026]
3. Woods J.D. and Watson N.R. 1970. "Measurements of Thermocline Fronts from the Air", Underwater Science & Technology Journal 2, 6, 90-99.
4. Oceanography of the Strait of Sicily, Proceedings of a Conference held at SACLANTCEN, 11-13 April 1972. SACLANTCEN Conference Proceedings CP-7, 1972.